

<https://helda.helsinki.fi>

---

## Prenatal Cotinine Levels and ADHD Among Offspring

Sourander, Andre

2019-03

---

Sourander , A , Sucksdorff , M , Chudal , R , Surcel , H-M , Hinkka-Yli-Salomaki , S ,  
Gyllenberg , D , Cheslack-Postava , K & Brown , A S 2019 , ' Prenatal Cotinine Levels and  
ADHD Among Offspring ' , Pediatrics (English Edition) , vol. 143 , no. 3 , 20183144 . <https://doi.org/10.1542/peds.2018-3144>

---

<http://hdl.handle.net/10138/313329>

<https://doi.org/10.1542/peds.2018-3144>

---

unspecified

publishedVersion

---

*Downloaded from Helda, University of Helsinki institutional repository.*

*This is an electronic reprint of the original article.*

*This reprint may differ from the original in pagination and typographic detail.*

*Please cite the original version.*

# Prenatal Cotinine Levels and ADHD Among Offspring

Andre Sourander, MD, PhD<sup>a</sup>, Minna Sucksdorff, MD<sup>b</sup>, Roshan Chudal, MBBS, MPH, PhD<sup>c</sup>, Heidi Marja Surcel, PhD,  
Susanna Hinkka-Yli-Salomäki, PhD<sup>d</sup>, David Gyllenberg, MD, PhD<sup>e</sup>, Deely Cheslack-Postava, PhD<sup>f</sup>, S. Brown, MD, MPH<sup>g</sup>

## abstract

**OBJECTIVE:** An association between maternal smoking during pregnancy and offspring attention-deficit/hyperactivity disorder (ADHD) has been shown across several studies based on self-reports. No previous studies have investigated the association of nicotine exposure measured by cotinine levels during pregnancy and offspring ADHD.

**METHODS:** In this population-based study, 1079 patients born between 1998 and 1999 and diagnosed with ADHD according to the International Classification of Diseases and 1079 matched controls were identified from Finnish nationwide registers. Maternal cotinine levels were measured by using quantitative immunoassays from maternal serum specimens collected during the first and second trimesters of pregnancy and archived in the national biobank.

**RESULTS:** There was a significant association between increasing log-transformed maternal cotinine levels and offspring ADHD. The odds ratio was 1.09 (95% confidence interval [CI] 1.06–1.12) when adjusting for maternal socioeconomic status, maternal age, maternal psychopathology, paternal age, paternal psychopathology, and child birth weight for gestational age. In the categorical analyses with cotinine levels in 3 groups, heavy nicotine exposure (cotinine level  $\geq 50$  ng/mL) was associated with offspring ADHD, with an odds ratio of 2.21 (95% CI 1.63–2.99) in the adjusted analyses. Analyses by deciles of cotinine levels revealed that the adjusted odds for offspring ADHD in the highest decile was 3.34 (95% CI 2.02–5.52).

**CONCLUSIONS:** The study reveals an association with and a dose-response relationship between nicotine exposure during pregnancy and offspring ADHD. Future studies incorporating maternal smoking and environmental, genetic, and epigenetic factors are warranted.



<sup>a</sup>Department of Child Psychiatry, University of Turku, Turku, Finland; <sup>b</sup>Department of Child Psychiatry and <sup>c</sup>Pediatric and Adolescent Medicine, Turku University Hospital, Turku, Finland; <sup>d</sup>New York State Psychiatric Institute, Department of Psychiatry, Columbia University Medical Center, New York, New York; <sup>e</sup>Medical Research Center, University of Oulu, Oulu, Finland; <sup>f</sup>Borealis of Northern Finland, Oulu University Hospital, Oulu, Finland; <sup>g</sup>National Institutes of Health and Welfare, Helsinki, Finland; <sup>h</sup>Department of Adolescent Psychiatry, Helsinki University Central Hospital, Helsinki, Finland; <sup>i</sup>Department of Epidemiology, Mailman School of Public Health, Columbia University, New York, New York

Dr Sourander conceptualized the study, participated in the study design, drafted the initial manuscript, and contributed to the interpretation of the data and the critical review and revision of the manuscript; Dr Sucksdorff participated in the study design, conducted the literature search, drafted portions of the initial manuscript, and contributed to the interpretation of the data and the critical review and revision of the manuscript; Dr Chudal participated in the study design, drafted portions of the initial manuscript, and contributed to the interpretation of the data and the critical review and revision of the manuscript; Drs Surcel, Gyllenberg, Cheslack-Postava, and Brown conceptualized the study, participated in the study design, and contributed to the interpretation of the data and the critical review and revision of the manuscript; Ms Hinkka-Yli-Salomäki designed the study, conducted the analyses, and contributed to the interpretation of the data and the critical review and revision of the manuscript; and all authors approved the manuscript as submitted.

**WHAT'S KNOWN ON THIS SUBJECT:** Exposure to maternal smoking is associated with various adverse perinatal outcomes. Association between maternal smoking and offspring attention-deficit/hyperactivity disorder has been shown across studies. However, the causality of the association has been questioned to be mostly due to familial confounding.

**WHAT THIS STUDY ADDS:** This first nationwide study, objectively measured nicotine exposure through maternal cotinine levels allows us to overcome underreporting of smoking during pregnancy. We report a strong association as well as a dose-dependent relationship between prenatal nicotine exposure and offspring attention-deficit/hyperactivity disorder.

**Key Words:** Sourander A, Sucksdorff M, Chudal R, et al. Prenatal Cotinine Levels and ADHD Among Offspring. *Pediatrics* 2019;143(3):e20183144











that quitting smoking was associated with less externalizing problems among offspring.<sup>38</sup> Furthermore, an American sibling study revealed that familial confounding explained inattentive but not hyperactive and/or impulsive ADHD behaviors, suggesting that the association may vary by phenotype.<sup>39</sup>

Despite several strengths, sibling comparison studies should also be interpreted with caution given their limitations. It has been suggested that within-sibling estimates will be biased toward the null by measurement error and that they may be either more or less biased than between-family estimates depending on the extent to which siblings share confounders versus the exposure.<sup>40</sup> In addition, mothers who vary in their smoking habits during different pregnancies can perhaps not be generalized to all smoking populations.<sup>40</sup>

It is possible that the association between nicotine exposure during pregnancy and ADHD might partially be explained also by gene-environment interaction.

Accumulating data provide evidence that prenatal smoking may act through epigenetic changes via altered DNA methylation and microRNA expression.<sup>41,42</sup> The exposure to nicotine may increase the risk of ADHD particularly among children with genetic vulnerability for ADHD. It is also possible that maternal smoking during pregnancy is a proxy risk factor leading to ADHD by independent mechanisms. Smoking during pregnancy is associated with poorer parenting skills, which are associated with child behavioral problems.<sup>43</sup>

The current study has several strengths, including being based on a large nationwide sample, assessing nicotine exposure with objective biological measurement and including several potential confounders. However, when interpreting the findings, several

limitations should be considered. A key question is whether smoking during pregnancy is causally associated with ADHD or is a proxy of another risk factor (eg, familial confounding). Even if most of the effect is due to familial or genetic confounding, the current study reveals that the association between smoking and ADHD has a dose-response effect. The limitation of observational data is that we cannot examine causal processes. We did not have access to biomarkers of sibling pregnancies that would have shed more light on a possible causal link between nicotine exposure during pregnancy and offspring ADHD. However, in the current study, we were able to adjust for several confounders, including BWGA, maternal SES, age, psychopathology, and substance use disorder as well as paternal psychopathology. Information about possible substance use during pregnancy was restricted to register-based substance use diagnoses. In the current study, 5.3% of mothers of children with ADHD had a diagnosis of substance use disorder, which was similar to the estimated prevalence in Finland of 6.4% for the exposure to maternal alcohol and drug dependence.<sup>44</sup> The number of parents diagnosed with ADHD in this sample was low, which is a limitation of the study (Supplemental Tables 5 and 6). The underdiagnoses among parents could be primarily because ADHD was not a widely used diagnosis in the parental generation. In addition, it is unlikely that ADHD among parents was treated in inpatient care. Because the FHDR did not cover outpatient diagnoses before 1998, the diagnosis of ADHD among parents is likely underestimated. Finally, the subjects with ADHD included in this study were only those who were referred to specialized services and likely represent the more severe cases of ADHD. However, authors of a previous study reported an 88% validity of the ADHD diagnoses in the FHDR

examined against Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition criteria for ADHD.<sup>7</sup>

## CONCLUSIONS

According to the World Health Organization, smoking is considered 1 of the main public health concerns worldwide.<sup>45</sup> The current study reveals a strong association between prenatal nicotine exposure and a dose-dependent relationship with offspring ADHD. This study adds 2 important aspects; first, the use of cotinine as a documented measure of nicotine exposure during pregnancy and second, the finding of a dose-response effect in the association. Given the high prevalence of both smoking during pregnancy and ADHD among children, these findings warrant future studies on the interplay between maternal smoking and environmental, genetic, and epigenetic factors.

## ACKNOWLEDGMENTS

We thank Mr Jesse Fomin, Ms Jarna Lindroos, and Mr Lauri Sillanmäki for their technical support.

## ABBREVIATIONS

ADHD: attention-deficit/hyperactivity disorder  
 BWGA: birth weight for gestational age  
 CI: confidence interval  
 FHDR: Finnish Hospital Discharge Register  
 FMBR: Finnish Medical Birth Register  
 FMC: Finnish Maternity Cohort  
 ICD-8: International Classification of Diseases, Eighth Revision  
 ICD-9: International Classification of Diseases, Ninth Revision  
 ICD-10: International Classification of Diseases, 10th Revision  
 OR: odds ratio  
 PIC: personal identity code  
 SES: socioeconomic status



## REFERENCES

- Drake P, Driscoll AK, Mathews TJ. Cigarette smoking during pregnancy: United States, 2011-2014. *NCHS Data Brief* 2018;(305):1-8.
- Heino A, Vuori E, Gissler M. Perinatal statistics - parturients, deliveries and newborns 2016 [in Finnish]. Available at: <http://urn.fi/URN:NBN:Fi-fe2017103150386>. Accessed August 15, 2018.
- Huang L, Wang Y, Zhang L, et al. Maternal smoking and attention deficit hyperactivity disorder in offspring: a meta-analysis. *Pediatrics*. 2018;141(1):e20172465.
- Tiesler CM, Heinrich J. Prenatal nicotine exposure and child behavioural problems. *Eur Child Adolesc Psychiatry* 2014;23(10):992-999.
- Obel C, Linnet KM, Henriksen TB, et al. Smoking during pregnancy and hyperactivity-inattention in the offspring: comparing results from three Nordic cohorts. *Int J Epidemiol* 2009;38(3):690-695.
- Langley K, Rice F, van den Bree MB, Thapar A. Maternal smoking during pregnancy as an environmental risk factor for attention deficit hyperactivity disorder behaviour. A review. *Dev Psychol* 2005;57(6):357-361.
- Joelsson P, Chudal R, Talati A, Suominen A, Brown AS, Sourander A. Prenatal smoking exposure and neuropsychiatric comorbidity of ADHD: a Finnish nationwide population-based cohort study. *BMC Psychiatry* 2016;16:306.
- Obel C, Zhu JL, Olsen J, et al. The risk of attention deficit hyperactivity disorder in children exposed to maternal smoking during pregnancy - a re-examination using a sibling design. *J Child Psychol Psychiatry* 2016;57(4):532-537.
- Skoglund C, Chen Q, D'Onofrio BM, Lichtenstein P, Larsson H. Familial confounding of the association between maternal smoking during pregnancy and ADHD in offspring. *Child Psychol Psychiatry* 2014;55(1):68-73.
- Gustavson K, Ystrom E, Stoltenberg C, et al. Smoking in pregnancy and child ADHD. *Pediatrics*. 2017;139(2):e20162509.
- Langley K, Heron J, Smith GD, Thapar A. Maternal and paternal smoking during pregnancy and risk of ADHD symptoms in offspring: testing for intrauterine effects. *Am J Epidemiol* 2012;176(3):261-268.
- D'Onofrio BM, Van Hulle CA, Waldman ID, et al. Smoking during pregnancy and offspring externalizing problems: an exploration of genetic and environmental confounding. *Dev Psychopathol* 2008;20(1):135-144.
- Lindblad F, Hjern A. ADHD after fetal exposure to maternal smoking. *Nicotine Tob Res*. 2010;12(4):408-413.
- Shipton D, Tappin DM, Vadiveloo T, Crossley JA, Aitken DA, Chalmers J. Reliability of self reported smoking status by pregnant women for estimating smoking prevalence: a retrospective, cross sectional study. *BMJ* 2009;339:b4347.
- Dietz PM, Homa D, England LJ, et al. Estimates of nondisclosure of cigarette smoking among pregnant and nonpregnant women of reproductive age in the United States. *Am J Epidemiol* 2011;173(3):355-359.
- Tong VT, Althabe F, Alemán A, et al; Prenatal Tobacco Cessation Intervention Collaborative. Accuracy of self-reported smoking cessation during pregnancy. *Acta Obstet Gynecol Scand* 2015;94(1):100-104.
- Connor Gorber S, Selhub H, Hurwitz S, Hardt J, Levasseur G, Tremblay M. The accuracy of self-reported smoking: a systematic review of the relationship between self-reported and cotinine-assessed smoking status. *Nicotine Tob Res*. 2009;11(1):12-22.
- Dürr DW, Høyer BB, Christensen LH, et al. Tobacco smoking during pregnancy and risk of adverse behaviour in offspring: a follow-up study. *Reprod Toxicol*. 2015;57:25-35.
- Eskenazi B, Trupin LS. Passive and active maternal smoking during pregnancy, as measured by serum cotinine, and postnatal smoke exposure. II. Effects on neurodevelopment at age 5 years. *Am J Epidemiol* 1995;142(suppl 9):S29-S39.

20. University of Oulu. Biobank Borealis of Northern Finland from bench to bedside. Available at: [www.oulu.university/node/38474](http://www.oulu.university/node/38474). Accessed March 14, 2018
21. Gissler M, Surcel H. Combining health register data and biobank data. *Stat J IAOS*. 2012;28(2):5358
22. World Health Organization. International Classification of Diseases, 10th Revision. Geneva, Switzerland: World Health Organization; 1992
23. World Health Organization. International Classification of Diseases, Ninth Revision. Geneva, Switzerland: World Health Organization; 1977
24. World Health Organization. International Classification of Diseases, Eighth Revision. Geneva, Switzerland: World Health Organization; 1967
25. Joelsson P, Chudal R, Gyllenberg D, et al. Demographic characteristics and psychiatric comorbidity of children and adolescents diagnosed with ADHD in specialized healthcare. *Child Psychiatry Hum Dev*. 2016;47(4):5582
26. Sankilampi U, Hannila ML, Saari A, Gissler M, Dunkel L. New population-based references for birth weight, length, and head circumference in singletons and twins from 23 to 43 gestation weeks. *Ann Med*. 2013;45(56):446454
27. Rothman KJ, Greenland S. *Modern Epidemiology*. 2nd ed. Philadelphia, PA: Lippincott-Raven; 1998
28. Niemelä S, Sourander A, Surcel HM, et al. Prenatal nicotine exposure and risk of schizophrenia among offspring in a national birth cohort. *Am J Psychiatry*. 2016;173(8):7996
29. Kapeu AS, Luostarinen T, Jellum E, et al. Is smoking an independent risk factor for invasive cervical cancer? A nested case-control study within Nordic biobanks. *Am J Epidemiol*. 2009;169(4):480488
30. Jauniaux E, Gulbis B, Acharya G, Thiry P, Rodeck C. Maternal tobacco exposure and cotinine levels in fetoids in the first half of pregnancy. *Obstet Gynecol*. 1999;93(1):279
31. Wickström R. Effects of nicotine during pregnancy: human and experimental evidence. *Curr Neuropharmacol*. 2007;5(3):213222
32. Bublitz MH, Stroud LR. Maternal smoking during pregnancy and offspring brain structure and function: review and agenda for future research. *Nicotine Tob Res*. 2012;14(4):398
33. Ekblad M, Korkeila J, Parkkola R, Lapinleimu H, Haataja L, Lehtonen L; PIPARI Study Group. Maternal smoking during pregnancy and regional brain volumes in preterm infants. *Pediatr*. 2010;156(2):1850.e1
34. Ekblad M, Korkeila J, Lehtonen L. Smoking during pregnancy affects foetal brain development. *Acta Paediatr*. 2015;104(1):113
35. Obel C, Olsen J, Henriksen TB, et al. Is maternal smoking during pregnancy a risk factor for hyperkinetic disorder? –Findings from a sibling design. *J Epidemiol*. 2011;40(2):335
36. Thapar A, Rice F, Hay D, et al. Prenatal smoking might not cause attention-deficit/hyperactivity disorder: evidence from a novel design. *Biol Psychiatry*. 2009;66(8):727
37. Ekblad M, Lehtonen L, Korkeila J, Gissler M. Maternal smoking during pregnancy and the risk of psychiatric morbidity in singleton sibling pairs. *Nicotine Tob Res*. 2017;19(5):5974
38. Dolan CV, Geels L, Vink JM, et al. Testing causal effects of maternal smoking during pregnancy on offspring externalizing and internalizing behavior. *Behav Genet*. 2016;46(3):378
39. Marceau K, Cinnamon Bidwell L, Karoly HC, et al. Within-family effects of smoking during pregnancy on ADHD: the importance of phenotype. *Abnorm Child Psychol*. 2018;46(4):659
40. Frisell T, Öberg S, Kuja-Halkola R, Sjölander A. Sibling comparison designs: bias from non-shared confounders and measurement error. *Epidemiology*. 2012;23(5):7720
41. Rzehak P, Saffery R, Reischl E, et al; European Childhood Obesity Trial Study group. Maternal smoking during pregnancy and DNA-methylation in children at age 5.5 years: epigenome-wide-analysis in the European Childhood Obesity Project (CHOP)-study. *PLoS One*. 2016;11(5):e0155554
42. Knopik VS, Maccani MA, Francrazio S, McGeary JE. The epigenetics of maternal cigarette smoking during pregnancy and effects on child development. *Dev Psychopathol*. 2012;24(4):13771390
43. Tandon M, Si X, Belden A, Spitznagel E, Wakschlag LS, Luby J. Parenting practices in pregnancy smokers compared to non smokers. *J Clin Med Res*. 2013;5(2):84
44. Pajulo M, Savonlahti E, Sourander A, Helenius H, Piha J. Antenatal depression, substance dependency and social support. *U Affect Disord*. 2001;65(1):917
45. World Health Organization. *MPower: A Policy Package to Reverse the Tobacco Epidemic*. Geneva, Switzerland: World Health Organization; 2008

## Prenatal Cotinine Levels and ADHD Among Offspring

Andre Sourander, Minna Sucksdorff, Roshan Chudal, Heljä-Marja Surcel, Susanna Hinkka-Yli-Salomäki, David Gyllenberg, Keely Cheslack-Postava and Alan Brown

*Pediatrics* 2019;143;

DOI: 10.1542/peds.2018-3144 originally published online February 25, 2019;

<b>Updated Information &amp; Services</b>	including high resolution figures, can be found at: <a href="http://pediatrics.aappublications.org/content/143/3/e20183144">http://pediatrics.aappublications.org/content/143/3/e20183144</a>
<b>References</b>	This article cites 38 articles, 3 of which you can access for free at: <a href="http://pediatrics.aappublications.org/content/143/3/e20183144#BIBL">http://pediatrics.aappublications.org/content/143/3/e20183144#BIBL</a>
<b>Subspecialty Collections</b>	This article, along with others on similar topics, appears in the following collection(s): <b>Developmental/Behavioral Pediatrics</b> <a href="http://www.aappublications.org/cgi/collection/development:behavioral_issues_sub">http://www.aappublications.org/cgi/collection/development:behavioral_issues_sub</a> <b>Attention-Deficit/Hyperactivity Disorder (ADHD)</b> <a href="http://www.aappublications.org/cgi/collection/attention-deficit:hyperactivity_disorder_adhd_sub">http://www.aappublications.org/cgi/collection/attention-deficit:hyperactivity_disorder_adhd_sub</a> <b>Substance Use</b> <a href="http://www.aappublications.org/cgi/collection/substance_abuse_sub">http://www.aappublications.org/cgi/collection/substance_abuse_sub</a> <b>Smoking</b> <a href="http://www.aappublications.org/cgi/collection/smoking_sub">http://www.aappublications.org/cgi/collection/smoking_sub</a>
<b>Permissions &amp; Licensing</b>	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: <a href="http://www.aappublications.org/site/misc/Permissions.xhtml">http://www.aappublications.org/site/misc/Permissions.xhtml</a>
<b>Reprints</b>	Information about ordering reprints can be found online: <a href="http://www.aappublications.org/site/misc/reprints.xhtml">http://www.aappublications.org/site/misc/reprints.xhtml</a>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



# PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

## **Prenatal Cotinine Levels and ADHD Among Offspring**

Andre Sourander, Minna Sucksdorff, Roshan Chudal, Heljä-Marja Surcel, Susanna Hinkka-Yli-Salomäki, David Gyllenberg, Keely Cheslack-Postava and Alan Brown  
*Pediatrics* 2019;143;

DOI: 10.1542/peds.2018-3144 originally published online February 25, 2019;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/143/3/e20183144>

Data Supplement at:

<http://pediatrics.aappublications.org/content/suppl/2019/02/20/peds.2018-3144.DCSupplemental>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2019 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

